

GROUP TOPOLOGIES COARSER THAN THE ISBELL TOPOLOGY

SZYMON DOLECKI, FRANCIS JORDAN AND FRÉDÉRIC MYNARD

ABSTRACT. The Isbell, compact-open and point-open topologies on the set $C(X, \mathbb{R})$ of continuous real-valued maps can be represented as the dual topologies with respect to some collections $\alpha(X)$ of compact families of open subsets of a topological space X . Those $\alpha(X)$ for which addition is jointly continuous at the zero function in $C_\alpha(X, \mathbb{R})$ are characterized, and sufficient conditions for translations to be continuous are found. As a result, collections $\alpha(X)$ for which $C_\alpha(X, \mathbb{R})$ is a topological vector space are defined canonically. The Isbell topology coincides with this vector space topology if and only if X is infraconsonant. Examples based on measure theoretic methods, that $C_\alpha(X, \mathbb{R})$ can be strictly finer than the compact-open topology, are given. To our knowledge, this is the first example of a splitting group topology strictly finer than the compact-open topology.

1. INTRODUCTION

The *Isbell*, *compact-open* and *point-open* topologies on the set $C(X, Y)$ of continuous real-valued maps from X to Y , can be represented as the *dual topologies* with regard to some collections $\alpha = \alpha(X)$ of compact openly isotone families of a topological space X , that is, the topology $\alpha(X, Y)$ is determined by a subbase of open sets of the form

$$(1.1) \quad [\mathcal{A}, U] := \{f \in C(X, \mathbb{R}) : f^-(U) \in \mathcal{A}\},$$

where $\emptyset \notin \mathcal{A} \in \alpha$ and U are open subsets of Y (and $f^-(U) := \{x \in X : f(x) \in U\}^1$). They are dual with regard to the collections, respectively, $\kappa(X)$ of all *compact families*, $k(X)$ of *compactly generated families* and $p(X)$ *finitely generated families* on X . In most cases, α is a topology on the space $\mathcal{O}_X$ of open subsets of X . The interrelations between the topology of X , the topology $\alpha(X)$, and the topology $\alpha(X, Y)$ can be, to a large extent, studied at this level of generality. In particular, under mild assumptions, local properties of $\alpha(X, \mathbb{R})$ at the zero function exactly correspond to local properties of $(\mathcal{O}_X, \alpha)$ at X . This greatly simplifies the study of local topological properties for the function spaces $C_\alpha(X, \mathbb{R})$, provided that $C_\alpha(X, \mathbb{R})$ is homogeneous. But while $p(X, \mathbb{R})$ and $k(X, \mathbb{R})$ are topological vector spaces for each X , the Isbell topology $\kappa(X, \mathbb{R})$ need not be even translation-invariant. In this context, it is natural to study under what conditions $\alpha(X, \mathbb{R})$ is translation-invariant, or even better, a topological group. If X is *consonant* (that is, if $k(X, \$^*) = \kappa(X, \$^*)$, where $\* designs the *Sierpiński topology*) then $k(X, \mathbb{R})$ and $\kappa(X, \mathbb{R})$ coincide, and in particular $\kappa(X, \mathbb{R})$ is a group topology. In [5] we characterized those topologies

Date: February 17, 2010.

¹ $f^-(U)$ is a shorthand for $f^{-1}(U)$

X , for which addition is jointly continuous at the zero function for the Isbell topology $\kappa(X, \mathbb{R})$; the class of such topologies, called *infraconsonant*, is larger than that of consonant topologies, but we do not know if the two classes coincide in case of completely regular topologies X . In this paper we prove that the Isbell topology $\kappa(X, \mathbb{R})$ is a group topology if and only if X is infraconsonant. Therefore the problem [5, Problem 1.1] of finding a completely regular infraconsonant space that is not consonant is equivalent to the problem of finding a completely regular space X such that the Isbell topology $\kappa(X, \mathbb{R})$ is a group topology, but does not coincide with the compact-open topology. While we do not solve this problem, we obtain the intermediate result that there are spaces X for which topologies $\alpha \subseteq \kappa(X)$ making $C_\alpha(X, \mathbb{R})$ a topological group strictly finer than $C_k(X, \mathbb{R})$ exist. Indeed, for each X there exists a largest hereditary ⁽²⁾ collection $\Lambda^\downarrow(X) \subseteq \kappa(X)$, for which the addition is jointly continuous at the zero function in $\Lambda^\downarrow(X, \mathbb{R})$. It turns out that $\Lambda^\downarrow(X, \mathbb{R})$ is a vector space topology and that a completely regular space X is infraconsonant if and only if $\Lambda^\downarrow(X, \mathbb{R}) = \kappa(X, \mathbb{R})$. Using measure theoretic methods, we show in particular that if a completely regular X is not pre-Radon, then $k(X, \mathbb{R})$ is strictly included in $\Lambda^\downarrow(X, \mathbb{R})$. Therefore each completely regular space X that is not pre-Radon admits a group topology on $C(X, \mathbb{R})$ that is splitting ⁽³⁾ but strictly finer than the compact-open topology. To our knowledge, no such example was known so far.

2. GENERALITIES

If $\mathcal{A}$ is a family of subsets of a topological space X then $\mathcal{O}_X(\mathcal{A})$ denotes the family of open subsets of X containing an element of $\mathcal{A}$. In particular, if $A \subset X$ then $\mathcal{O}_X(A)$ denotes the family of open subsets of X containing A . We denote by $\mathcal{O}_X$ the set of open subsets of X .

If X and Y are topological spaces, $C(X, Y)$ denotes the set of continuous functions from X to Y . If $A \subset X$, $U \subset Y$, then $[A, U] := \{f \in C(X, Y) : f(A) \subset U\}$. A family $\mathcal{A}$ of subsets of X is *openly isotone* if $\mathcal{O}_X(\mathcal{A}) = \mathcal{A}$. If $\mathcal{A}$ is openly isotone and U is open, then $[\mathcal{A}, U] = \bigcup_{A \in \mathcal{A}} [A, U]$.

If α is a collection of openly isotone families $\mathcal{A}$ of open subsets of X , such that each open subset of X belongs to an element of α , then

$$\{[\mathcal{A}, U] : \mathcal{A} \in \alpha, U \in \mathcal{O}_Y\}$$

forms a subbase for a topology $\alpha(X, Y)$ on $C(X, Y)$. We denote the set $C(X, Y)$ endowed with this topology by $C_\alpha(X, Y)$. Note that because

$$[\mathcal{A}, U] \cap [\mathcal{B}, U] = [\mathcal{A} \cap \mathcal{B}, U],$$

$\alpha(X, Y)$ and $\alpha^\cap(X, Y)$ coincide, where $\alpha^\cap$ consists of finite intersections of the elements of α . Therefore, we can always assume that α is stable under finite intersections.

In the sequel, we will focus on the case where α consists of compact families. A family $\mathcal{A} = \mathcal{O}_X(\mathcal{A})$ is *compact* if whenever $\mathcal{P} \subset \mathcal{O}_X$ and $\bigcup \mathcal{P} \in \mathcal{A}$ then there is a

²A collection α is *hereditary* if $\mathcal{A} \downarrow A \in \alpha$ whenever $A \in \mathcal{A} \in \alpha$, where $\mathcal{A} \downarrow A$ is defined by (2.2).

³that is, if for each space Y , the continuity of a map $g : Y \times X \rightarrow \mathbb{R}$ implies the continuity of the map $g^* : Y \rightarrow C_\alpha(X, \mathbb{R})$ defined by $g^*(y)(x) = g(y, x)$; equivalently, a topology is splitting if it is coarser than the continuous convergence.

finite subfamily $\mathcal{P}_0$ of $\mathcal{P}$ such that $\bigcup \mathcal{P}_0 \in \mathcal{A}$. Of course, for each compact subset K of X , the family $\mathcal{O}_X(K)$ is compact.

We denote by $\kappa(X)$ the collection of compact families on X . Seen as a family of subsets of $\mathcal{O}_X$ (the set of open subsets of X), $\kappa(X)$ is the set of open sets for the *Scott topology*; hence every union of compact families is compact, in particular $\bigcup_{K \in \mathcal{K}} \mathcal{O}_X(K)$ is compact if $\mathcal{K}$ is a family of compact subsets of X . A topological space is called *consonant* if every compact family $\mathcal{A}$ is *compactly generated*, that is, there is a family $\mathcal{K}$ of compact sets such that $\mathcal{A} = \bigcup_{K \in \mathcal{K}} \mathcal{O}_X(K)$. Similarly, $p(X) := \{\mathcal{O}_X(F) : F \in [X]^{<\omega}\}$ and $k(X) := \{\mathcal{O}(K) : K \subseteq X \text{ compact}\}$ are basis for topologies on $\mathcal{O}_X$. Accordingly, $p(X, Y)$ is the *topology of pointwise convergence*, $k(X, Y)$ is the *compact-open topology* and $\kappa(X, Y)$ is the *Isbell topology* on $C(X, Y)$.

If $\mathbb{S}^* := \{\emptyset, \{0\}, \{0, 1\}\}$ the function spaces $C(X, \mathbb{S}^*)$ can be identified with the set of open subsets of X ⁽⁴⁾. In this notation, X is consonant if and only if $C_k(X, \mathbb{S}^*) = C_\kappa(X, \mathbb{S}^*)$.

More generally, a space X is called *Z-consonant* if $C_\kappa(X, Z) = C_k(X, Z)$ [11, chapter 3]. [11, Problem 62] asks for what spaces Z (other than $\mathbb{S}^*$) Z -consonance implies consonance. A still more general problem is, given a collection α of compact families defined for each space X , to determine for what spaces Z ,

$$(2.1) \quad C_k(X, Z) = C_\alpha(X, Z) \iff C_k(X, \mathbb{S}^*) = C_\alpha(X, \mathbb{S}^*).$$

The latter equality always implies the former. More generally, in view of the definition of $\alpha(X, Z)$, if α and γ are collections (of compact families on X), then

$$C_\alpha(X, \mathbb{S}^*) \leq C_\gamma(X, \mathbb{S}^*) \implies C_\alpha(X, Z) \leq C_\gamma(X, Z)$$

for every topological space Z . To show the converse implication under some additional assumptions, recall that the *restriction of $\mathcal{A}$ to $A \in \mathcal{A}$* is defined by

$$(2.2) \quad \mathcal{A} \downarrow A := \{U \in \mathcal{O}_X : \exists B \subseteq A \cap U, B \in \mathcal{A}\}.$$

[5, Lemma 2.8] shows that if $\mathcal{A}$ is a compact family and $A \in \mathcal{A}$, then $\mathcal{A} \downarrow A$ is compact too. A collection α of families of open subsets of a given set is *hereditary* if $\mathcal{A} \downarrow A \in \alpha$ whenever $\mathcal{A} \in \alpha$ and $A \in \mathcal{A}$.

It was shown in [5, Proposition 2.4] that if X is completely regular and $\mathbb{R}$ -consonant, then it is consonant. More generally:

Proposition 2.1. *If $\alpha, \gamma \subseteq \kappa(X)$ are two topologies, α is hereditary, X is completely regular, and $C_\alpha(X, \mathbb{R}) \leq C_\gamma(X, \mathbb{R})$, then $C_\alpha(X, \mathbb{S}^*) \leq C_\gamma(X, \mathbb{S}^*)$.*

Proof. The neighborhood filter of an open set A with respect to $\alpha(X, \mathbb{S}^*)$ is generated by a base of the form $\{\mathcal{A} \in \alpha : A \in \mathcal{A}\}$. Therefore we need show that for each $\mathcal{A} \in \alpha$ and each $A \in \mathcal{A}$, there exists $\mathcal{G} \in \gamma$ such that $\mathcal{G} \subseteq \mathcal{A} \downarrow A$. By assumption, $\mathcal{N}_\gamma(\overline{0}) \geq \mathcal{N}_\alpha(\overline{0})$ so that for each $\mathcal{A} \in \alpha$ and each $A \in \mathcal{A}$, there exists $\mathcal{G} \in \gamma$ and $r > 0$ such that $[\mathcal{G}, (-r, r)] \subset [\mathcal{A} \downarrow A, (-\frac{1}{2}, \frac{1}{2})]$. Suppose that there exists $G \in \mathcal{G} \setminus (\mathcal{A} \downarrow A)$, hence $X \setminus G \in (\mathcal{A} \downarrow A)^\#$. Because X is completely regular and $\mathcal{G}$ is compact there is $G_0 \in \mathcal{G}$ and a continuous function f such that $f(G_0) = \{0\}$ and $f(X \setminus G) = \{1\}$, by [5, Lemma 2.5]. Then $f \in [\mathcal{G}, (-r, r)]$ but $f \notin [\mathcal{A} \downarrow A, (-\frac{1}{2}, \frac{1}{2})]$,

⁴In [3], [5] and [6], we distinguish two homeomorphic copies $\mathbb{S} := \{\emptyset, \{1\}, \{0, 1\}\}$ and $\mathbb{S}^* := \{\emptyset, \{0\}, \{0, 1\}\}$ of the *Sierpiński topology* on $\{0, 1\}$ and identify the function spaces $C(X, \mathbb{S})$ and $C(X, \mathbb{S}^*)$ with the set of closed subsets of X and open subsets of X respectively. This is why we use $\mathbb{S}^*$ here.

because $1 \in f(B)$ for each $B \in \mathcal{A} \downarrow A$. Therefore $A \in \mathcal{G} \subseteq \mathcal{A} \downarrow A \subseteq \mathcal{A}$, so that $\alpha \leq \gamma$. $\square$

Corollary 2.2. *If X is completely regular and $\alpha \subseteq \kappa(X)$ is hereditary, then (2.1) holds for $Z = \mathbb{R}$.*

The *grill* of a family $\mathcal{A}$ of subsets of X is the family $\mathcal{A}^\# := \{B \subseteq X : \forall A \in \mathcal{A}, A \cap B \neq \emptyset\}$. Note that if $\mathcal{A} = \mathcal{O}(\mathcal{A})$, then

$$A \in \mathcal{A} \iff A^c \notin \mathcal{A}^\#.$$

If $\mathcal{A} \in \kappa(X)$ and C is a closed subset of X such that $C \in \mathcal{A}^\#$ then the family

$$\mathcal{A} \vee C := \mathcal{O}(\{A \cap C : A \in \mathcal{A}\}),$$

called *section of $\mathcal{A}$ by C* , is a compact family on X [3]. A collection α of families of open subsets of a given set is *sectionable* if $\mathcal{A} \vee C \in \alpha$ whenever $\mathcal{A} \in \alpha$ and C is a closed set in $\mathcal{A}^\#$. It was shown in [5, Theorem 2.9] that $C_\kappa(X, Z)$ is completely regular whenever Z is. A simple modification of the proof leads to the following generalization.

Theorem 2.3. *If Z is completely regular and $\alpha \subseteq \kappa(X)$ is sectionable, then $C_\alpha(X, Z)$ is completely regular.*

As $r[\mathcal{A}, U] = [\mathcal{A}, rU]$ for all $r \neq 0$, it is immediate that inversion for $+$ is always continuous in $C_\alpha(X, \mathbb{R})$. More generally, the proof of the joint continuity of scalar multiplication in $C_\kappa(X, \mathbb{R})$ [5, Proposition 2.10] can be adapted to the effect that:

Proposition 2.4. *If $\alpha \subseteq \kappa(X)$ is hereditary, then multiplication by scalars is jointly continuous for $C_\alpha(X, \mathbb{R})$.*

Corollary 2.5. *Let $\alpha \subseteq \kappa(X)$ be hereditary. If $C_\alpha(X, \mathbb{R})$ is a topological group then it is a topological vector space.*

3. SELF-JOINABLE COLLECTIONS AND JOINT CONTINUITY OF ADDITION AT THE ZERO FUNCTION

As usual, if A and B are subsets of an additive group, $A + B := \{a + b : a \in A, b \in B\}$ and if $\mathcal{A}$ and $\mathcal{B}$ are two families of subsets, $\mathcal{A} + \mathcal{B} := \{A + B : A \in \mathcal{A}, B \in \mathcal{B}\}$.

As we have mentioned, a topology on an additive group is a group topology if and only if inversion and translations are continuous, and $\mathcal{N}(o) + \mathcal{N}(o) \geq \mathcal{N}(o)$, where o is the neutral element. First, we investigate the latter property, that is,

$$(3.1) \quad \mathcal{N}_\alpha(\overline{0}) + \mathcal{N}_\alpha(\overline{0}) \geq \mathcal{N}_\alpha(\overline{0}),$$

for the space $C_\alpha(X, \mathbb{R})$, where $\overline{0}$ denotes the zero function.

If α and γ are two subsets of $\kappa(X)$, we say that α is γ -joinable if for every $\mathcal{A} \in \alpha$, there is $\mathcal{G} \in \gamma$ such that $\mathcal{G} \vee \mathcal{G} \subseteq \mathcal{A}$, where

$$\mathcal{G} \vee \mathcal{G} := \{G_1 \cap G_2 : G_1, G_2 \in \mathcal{G}\}.$$

A subset α of $\kappa(X)$ is *self-joinable* if it is α -joinable. A family $\mathcal{A}$ is called *joinable* if $\{\mathcal{A}\}$ is $\kappa(X)$ -joinable. In [5], a space X is called *infraconsonant* if every compact family is joinable, that is, $\kappa(X)$ is self-joinable. [5, Theorem 3.1] shows that among completely regular spaces X ,

$$\mathcal{N}_\kappa(\overline{0}) + \mathcal{N}_\kappa(\overline{0}) \geq \mathcal{N}_\kappa(\overline{0})$$

if and only if X is infraconsonant. More generally,

Theorem 3.1. *Let X be a completely regular space. Then $\alpha \subseteq \kappa(X)$ is self-joinable if and only if*

$$(3.2) \quad \mathcal{N}_\alpha(\overline{0}) + \mathcal{N}_\alpha(\overline{0}) \geq \mathcal{N}_\alpha(\overline{0}).$$

Proof. Let $\mathcal{A} \in \alpha$ and $V \in \mathcal{N}_\mathbb{R}(0)$. Because α is self-joinable, there exist a compact family $\mathcal{B}$ in α such that $\mathcal{B} \vee \mathcal{B} \subseteq \mathcal{A}$. If $W \in \mathcal{N}_\mathbb{R}(0)$ such that $W + W \subseteq V$, then $[\mathcal{B}, W] + [\mathcal{B}, W] \subseteq [\mathcal{A}, V]$, which proves (3.2).

Conversely, assume that $\alpha = \alpha^\cap$ is not self-joinable. Let $\mathcal{A}$ be a family of α such that $\mathcal{B} \vee \mathcal{B} \not\subseteq \mathcal{A}$ for every $\mathcal{B} \in \alpha$. Note that $\mathcal{B} \vee \mathcal{C} \not\subseteq \mathcal{A}$ for every pair of families $\mathcal{B}$ and $\mathcal{C}$ in α for otherwise $\mathcal{D} = \mathcal{B} \cap \mathcal{C}$ would be a family of α such that $\mathcal{D} \vee \mathcal{D} \subseteq \mathcal{A}$. Let $V = (-\frac{1}{2}, \frac{1}{2})$. We claim that for any pair $(\mathcal{B}, \mathcal{C}) \in \alpha^2$ and any pair (U, W) of $\mathbb{R}$ -neighborhood of 0, $[\mathcal{B}, U] + [\mathcal{C}, W] \not\subseteq [\mathcal{A}, V]$. Indeed, there exist $B \in \mathcal{B}$ and $C \in \mathcal{C}$ such that $B \cap C \notin \mathcal{A}$. Then $B^c \cup C^c \in \mathcal{A}^\#$. Moreover, $B^c \notin \mathcal{B}^\#$ so that by [5, Lemma 2.5], there exist $B_1 \in \mathcal{B}$ and $f \in C(X, \mathbb{R})$ such that $f(B_1) = \{0\}$ and $f(B^c) = \{1\}$. Similarly, $C^c \notin \mathcal{C}$ so that there exist $C_1 \in \mathcal{C}$ and $g \in C(X, \mathbb{R})$ such that $g(C_1) = \{0\}$ and $g(C^c) = \{1\}$. Then $f + g \in [\mathcal{B}, U] + [\mathcal{C}, W]$ but $1 \in (f + g)(A)$ for all $A \in \mathcal{A}$ so that $f + g \notin [\mathcal{A}, V]$. $\square$

Note that the collection $k(X)$ (of compactly generated families) is self-joinable, and that a union of self-joinable collections is self-joinable. Therefore, there is a largest self-joinable subset $\Lambda(X)$ of $\kappa(X)$. If α is self-joinable, so is $\alpha^\cap$. Therefore $\Lambda(X)$ is stable for finite intersections. In fact, $\Lambda(X)$ is a topology on $C(X, \mathbb{R}^*)$ and

$$(3.3) \quad k(X) \subseteq \Lambda(X) \subseteq \kappa(X).$$

Corollary 3.2. *Let X be completely regular. The largest subcollection α of $\kappa(X)$, for which (3.2) holds is $\alpha = \Lambda(X)$. In particular, a completely regular space X is infraconsonant if and only if $\kappa(X) = \Lambda(X)$.*

The collection $\Lambda(X)$ is sectionable ⁽⁵⁾, but in general it is not hereditary. We will construct now a largest hereditary collection of compact families for which (3.2) holds.

If α and γ are two subsets of $\kappa(X)$, we say that α is *hereditarily γ -joinable* if for every $\mathcal{A} \in \alpha$, and every $A \in \mathcal{A}$, there is $\mathcal{G} \in \gamma$ such that $A \in \mathcal{G}$ and $\mathcal{G} \vee \mathcal{G} \subseteq \mathcal{A}$. A subset α of $\kappa(X)$ is *hereditarily self-joinable* if it is hereditarily α -joinable. A family $\mathcal{A}$ is called *hereditarily joinable* if $\{\mathcal{A}\}$ is hereditarily $\kappa(X)$ -joinable. There exists a largest hereditarily self-joinable subset $\Lambda^\downarrow(X)$ of $\kappa(X)$. Notice that $\Lambda^\downarrow(X)$ is also the largest self-joinable and hereditary collection of compact families, and that $\Lambda^\downarrow(X)$ is sectionable.

Corollary 3.3. *Let X be completely regular. The largest hereditary subcollection α of $\kappa(X)$, for which (3.2) holds is $\alpha = \Lambda^\downarrow(X)$. In particular, a completely regular space X is infraconsonant if and only if $\kappa(X) = \Lambda^\downarrow(X)$.*

Of course, $\Lambda^\downarrow(X)$ is a topology, and $\Lambda^\downarrow(X) \subseteq \Lambda(X)$. The inclusion can be strict. In fact, we have:

Proposition 3.4. *A regular space X is infraconsonant if and only if $\Lambda^\downarrow(X) = \Lambda(X)$ if and only if $\kappa(X) = \Lambda^\downarrow(X)$.*

⁵Indeed, if α is self-joinable, so is $\{\mathcal{A} \vee \mathcal{C} : \mathcal{A} \in \alpha, \mathcal{C} \in \mathcal{A}^\#, \mathcal{C} \text{ closed}\}$ because if $\mathcal{A} \in \alpha$, there is $\mathcal{B} \in \alpha$ such that $\mathcal{B} \vee \mathcal{B} \subseteq \mathcal{A} \subseteq \mathcal{A} \vee \mathcal{C}$. By maximality, $\Lambda(X)$ is sectionable.

Proof. If X is not infraconsonant, there is a non-joinable family $\mathcal{A}$ on X . For any $x \in X \setminus \bigcap \mathcal{A}$, the family $\mathcal{O}(x) \cup \mathcal{A}$ belongs to $\Lambda(X)$ but not to $\Lambda^\downarrow(X)$. If X is regular and infraconsonant, then by [5, Lemma 3.2] $\kappa(X) = \Lambda^\downarrow(X)$. Finally if $\kappa(X) = \Lambda^\downarrow(X)$ then $\Lambda^\downarrow(X) = \Lambda(X)$, because $\Lambda(X)$ is between $\Lambda^\downarrow(X)$ and $\kappa(X)$. $\square$

Examples of non-infraconsonant spaces are provided in [5], so that both inclusions in (3.3) can be strict simultaneously.

4. SELF-SPLITTABLE COLLECTIONS AND CONTINUITY OF TRANSLATIONS

A collection α is γ -*splittable* if for every $\mathcal{A} \in \alpha$ and for every open subsets U_1 and U_2 of X such that $U_1 \cup U_2 \in \mathcal{A}$ there exist families $\mathcal{G}_i = \mathcal{G}_i \downarrow U_i$ in γ , $i \in \{1, 2\}$, such that $\mathcal{G}_1 \cap \mathcal{G}_2 \subseteq \mathcal{A}$. A collection α is *self-splittable* if it is α -splittable. A compact family $\mathcal{A}$ on X is *splittable* if $\{\mathcal{A}\}$ is $\kappa(X)$ -splittable. An immediate induction shows that if α is self-splittable, then for every $\mathcal{A} \in \alpha$ and every finite collection $\{U_1, \dots, U_n\}$ of open subsets of X such that $\bigcup_{i=1}^n U_i \in \mathcal{A}$, there are families $\mathcal{C}_i \in \alpha$ with $\mathcal{C}_i = \mathcal{C}_i \downarrow U_i$ such that $\bigcap_{i=1}^n \mathcal{C}_i \subseteq \mathcal{A}$.

In [10], F. Jordan calls a topological space *compactly splittable* if every compact family is splittable. A topological space with at most one non-isolated point is said to be *prime*. A modification of the proof of [9, Theorem 18] shows:

Proposition 4.1. *Prime spaces are compactly splittable.*

It follows from [10, Theorem 2] and [5, Corollary 4.2] that translations are continuous in the Isbell topological space $C_\kappa(X, \mathbb{R})$ if X is compactly splittable. More generally, we have:

Proposition 4.2. *If $\alpha \subseteq \kappa(X)$ is self-splittable, then translations are continuous for $C_\alpha(X, \mathbb{R})$.*

Proof. We show continuity of the translation by f_0 at g_0 . Let $\mathcal{A} \in \alpha$ and $U \in \mathcal{O}_\mathbb{R}$ such that $f_0 + g_0 \in [\mathcal{A}, U]$. There is $A_0 \in \mathcal{A}$ such that $(f_0 + g_0)(A_0) \subseteq U$. For each $x \in A_0$, there exists $V_x = -V_x \in \mathcal{O}_\mathbb{R}(0)$ such that $f_0(x) + g_0(x) + 2V_x \subseteq U$. Moreover, by continuity of f_0 and g_0 , there is $W_x \in \mathcal{O}_X(x)$ such that $f_0(W_x) \subseteq f_0(x) + V_x$ and $g_0(W_x) \subseteq g_0(x) + V_x$. As $\bigcup_{x \in A_0} W_x \in \mathcal{A}$ and $\mathcal{A}$ is compact, there is a finite subset F of A_0 such that $W := \bigcup_{x \in F} W_x \in \mathcal{A}$. Because α is self-splittable, there exists, for each $x \in F$, a compact family $\mathcal{C}_x = \mathcal{C}_x \downarrow W_x$ of α such that $\bigcap_{x \in F} \mathcal{C}_x \subseteq \mathcal{A}$. Note that by construction $g_0 \in \bigcap_{x \in F} [\mathcal{C}_x, g_0(x) + V_x]$. Moreover, if $g \in \bigcap_{x \in F} [\mathcal{C}_x, g_0(x) + V_x]$, then for each $x \in F$ there is $C_x \in \mathcal{C}_x$, $C_x \subseteq W_x$, such that $g(C_x) \subseteq g_0(x) + V_x$. If $y \in \bigcup_{x \in F} C_x \in \mathcal{A}$, then $y \in C_x$ for some x and

$$f_0(y) + g(y) \in f_0(x) + V_x + g_0(x) + V_x \subseteq U,$$

so that $f_0 + \bigcap_{x \in F} [\mathcal{C}_x, g_0(x) + V_x] \subseteq [\mathcal{A}, U]$. $\square$

[3, Example 4.9] shows that even on a compactly splittable space (like the Arens space), there exists $\alpha \subseteq \kappa(X)$ such that translations are not continuous for $C_\alpha(X, \mathbb{R})$, so that α is not self-splittable.

Note that in a regular space X , the collection $k(X)$ is self-splittable, and a union of self-splittable collections is self-splittable. Therefore, there is a largest self-splittable subset $\Sigma(X)$ of $\kappa(X)$. If α is self-splittable, so is $\alpha^\cap$, hence that $\Sigma(X)$ is a topology on $C(X, \mathbb{R}^*)$, and

$$(4.1) \quad k(X) \subseteq \Sigma(X) \subseteq \kappa(X).$$

Moreover, $\Sigma(X)$ is clearly hereditary, and sectionable ⁽⁶⁾.

Both inequalities in (4.1) can be strict. Examples of non-compactly splittable spaces are provided in [9], so that $\Sigma(X)$ can be strictly included in $\kappa(X)$. On the other hand, in view of Proposition 4.1, if X is prime and not consonant (e.g., the Arens space), then $k(X)$ is strictly included in $\Sigma(X)$.

Theorem 4.3. *Let X be regular. If $\alpha \subseteq \kappa(X)$ is self-joinable, hereditary, and sectionable, then α is self-splittable.*

Proof. By way of contradiction, assume that α is not self-splittable. Let $\alpha_1 = \alpha \cup \{\mathcal{O}(X)\}$. Using that $\emptyset \in \mathcal{O}(X)$, one can easily show that α_1 is self-joinable, hereditary, and sectionable. It is also easy to check that α_1 is not self-splittable.

Let $\mathcal{C} \in \alpha_1$ witness that α_1 is not self-splittable. There exist $\mathcal{C} \in \alpha_1$ and open sets U_1, U_2 such that $U_1 \cup U_2 \in \mathcal{C}$, but for any $\mathcal{B}_1, \mathcal{B}_2 \in \alpha_1$ with $U_1 \in \mathcal{B}_1$ and $U_2 \in \mathcal{B}_2$ we have $\mathcal{B}_1 \cap \mathcal{B}_2 \notin \mathcal{C}$.

By regularity, there is an open cover $\mathcal{V}$ of $U_1 \cup U_2$ such that $\text{cl}(V) \subseteq U_1$ or $\text{cl}(V) \subseteq U_2$ for every $V \in \mathcal{V}$. Since $U_1 \cup U_2 \in \mathcal{C}$, there exist a finite $\mathcal{V}_1 \subseteq \mathcal{V}$ such that $\bigcup \mathcal{V}_1 \in \mathcal{C}$. Let $W_1 = \bigcup \{V \in \mathcal{V}_1 : \text{cl}(V) \subseteq U_1\}$ and $W_2 = \bigcup \{V \in \mathcal{V}_1 : \text{cl}(V) \subseteq U_2\}$. Notice that $W_1 \cup W_2 \in \mathcal{C}$, $\text{cl}(W_1) \subseteq U_1$, and $\text{cl}(W_2) \subseteq U_2$. Let $\mathcal{C}_1 = \mathcal{C} \downarrow (W_1 \cup W_2)$. Since α_1 is hereditary, $\mathcal{C}_1 \in \alpha_1$. By self-joinability of α_1 there is a $\mathcal{D} \in \alpha_1$ such that $\mathcal{D} \vee \mathcal{D} \subseteq \mathcal{C}_1$. Notice that $\mathcal{D} \subseteq \mathcal{C}_1$.

Suppose there is a $D \in \mathcal{D}$ such that $D \cap W_1 = \emptyset$. Since $D \in \mathcal{D} \subseteq \mathcal{C}_1$, there is an $E \in \mathcal{C}$ such that $E \subseteq D \cap (W_1 \cup W_2)$. Notice that $E \subseteq W_2 \subseteq U_2$. So, $U_2 \in \mathcal{C}$. Since α_1 is hereditary, $\mathcal{C} \downarrow U_2 \in \alpha_1$. Notice that $U_2 \in \mathcal{C} \downarrow U_2 \in \alpha_1$, $U_1 \in \mathcal{O}(X) \in \alpha_1$, and $\mathcal{C} \downarrow U_2 \cap \mathcal{O}(X) = \mathcal{C} \downarrow U_2 \subseteq \mathcal{C}$, which contradicts our choice of U_1 and U_2 . So, we may assume that $\mathcal{D} \# W_1$. Similarly, we may assume that $\mathcal{D} \# W_2$.

For each $i \in \{1, 2\}$ let $\mathcal{D}_i = \mathcal{D} \vee \text{cl}(W_i)$. Since α_1 is sectionable, $\mathcal{D}_1, \mathcal{D}_2 \in \alpha_1$. Notice that $U_i \in \mathcal{D}_i$ for every i . Let $P \in \mathcal{D}_1 \cap \mathcal{D}_2$. For every $i \in \{1, 2\}$ there exist $D_i \in \mathcal{D}$ such that $D_i \cap \text{cl}(W_i) \subseteq P$. Since $D_1, D_2 \in \mathcal{D}$, $D_1 \cap D_2 \in \mathcal{C}_1$. There is an $E \in \mathcal{C}$ such that $E \subseteq D_1 \cap D_2 \cap (W_1 \cup W_2)$. Now,

$$E \subseteq (D_1 \cap D_2) \cap \text{cl}(W_1 \cup W_2) \subseteq (D_1 \cap \text{cl}(W_1)) \cup (D_2 \cap \text{cl}(W_2)) \subseteq P.$$

So, $P \in \mathcal{C}$. Thus, $\mathcal{D}_1 \cap \mathcal{D}_2 \subseteq \mathcal{C}$, contradicting our choice of U_1 and U_2 . $\square$

Corollary 4.4. *Let X be regular. If X is infraconsonant then X is compactly splittable.*

Proof. If X is infraconsonant, then $\kappa(X)$ is self-joinable. Since $\kappa(X)$ is hereditary and sectionable, $\kappa(X)$ is self-splittable, by Theorem 4.3. Since $\kappa(X)$ is self-splittable, X is compactly splittable. $\square$


Note that Corollary 4.4 provides a negative answer to [5, Problem 1.2]. The converse of Corollary 4.4 is not true. For instance, the Arens space is compactly splittable because it is prime, but it is not infraconsonant [5, Theorem 3.6].

⁶Indeed, if α is self-splittable, so is $\{\mathcal{A} \vee \mathcal{C} : \mathcal{A} \in \alpha, \mathcal{C} \in \mathcal{A}^\#, \mathcal{C} \text{ closed}\}$. Indeed, if $\bigcup_{i=1}^n U_i \in \mathcal{A} \vee \mathcal{C}$, then $\bigcup_{i=1}^n U_i \cup \mathcal{C}^c \in \mathcal{A}$, so that there are families $\mathcal{C}_i = \mathcal{C}_i \downarrow U_i$ and $\mathcal{C}_c = \mathcal{C}_c \downarrow \mathcal{C}^c$ in α such that $\bigcap \mathcal{C}_i \cap \mathcal{C}_c \subseteq \mathcal{A}$. Therefore

$$\bigcap_i \mathcal{C}_i \vee \mathcal{C} = \left(\bigcap_i \mathcal{C}_i \cap \mathcal{C}_c \right) \vee \mathcal{C} \subseteq \mathcal{A} \vee \mathcal{C}.$$

By maximality, $\Lambda(X)$ is sectionable.

In the diagram below, X is a regular space, and arrows represent inclusions. We have already justified that all of these inclusions may be strict, except for $k(X) \subseteq \Lambda^\downarrow(X)$. We will see in the next section that it may be strict.



In view of Proposition 2.4, Theorem 3.1, Proposition 4.2 and Theorem 4.3, we obtain:

Corollary 4.5. *Let X be completely regular. Then $C_{\Lambda^\downarrow}(X, \mathbb{R})$ is a topological vector space.*

As a consequence, we can extend [5, Theorem 5.3] from prime spaces to general completely regular spaces to the effect that:

Corollary 4.6. *Let X be completely regular. The following are equivalent:*

- (1) X is infraconsonant;
- (2) $\kappa(X) = \Lambda(X)$;
- (3) $\kappa(X) = \Lambda^\downarrow(X)$;
- (4) $\Lambda(X) = \Lambda^\downarrow(X)$;
- (5) $C_\kappa(X, \mathbb{R})$ is a topological vector space;
- (6) $C_\kappa(X, \mathbb{R})$ is a topological group;
- (7) $\mathcal{N}_\kappa(\overline{0}) + \mathcal{N}_\kappa(\overline{0}) \geq \mathcal{N}_\kappa(\overline{0})$;
- (8) $\cap : C_\kappa(X, \$^*) \times C_\kappa(X, \$^*) \rightarrow C_\kappa(X, \$^*)$ is jointly continuous.

Proof. Equivalences between (1) through (7) follow immediately from Theorem 3.1, Corollary 3.2 and Proposition 3.4. The equivalence with (8) follows from [5, Proposition 3.3]. $\square$

5. A VECTOR SPACE TOPOLOGY STRICTLY FINER THAN THE COMPACT-OPEN TOPOLOGY

A finite measure μ is called τ -additive if for every family $\mathcal{P} \subset \mathcal{O}_X$, and for every $\varepsilon > 0$ there is a finite subfamily $\mathcal{P}_\varepsilon \subset \mathcal{P}$ such that $\mu(\bigcup \mathcal{P}_\varepsilon) \geq \mu(\bigcup \mathcal{P}) - \varepsilon$. Hence, if μ is a τ -additive measure on X , then for each $r > 0$, the family

$$\mathcal{M}_r^\mu := \{O \in \mathcal{O}_X : \mu(O) > r\}$$

is compact. A topological space X is called *pre-Radon* if every finite τ -additive measure μ on X is a Radon measure, that is, $\mu(B) = \sup \{\mu(K) : K \subset B, K \text{ compact}\}$ for each Borel subset B of X .

Lemma 5.1. *Let μ be a τ -additive finite measure on a space X . Then $\gamma_\mu := \{\mathcal{M}_r^\mu \downarrow A : A \in \mathcal{M}_r^\mu, r > 0\}$ is self-splittable and hereditarily self-joinable.*

Proof. *Proof of 1.* γ_μ is hereditary and self-joinable (hence hereditarily self-joinable) because if $U \in \mathcal{M}_r^\mu$ and $m = \frac{r + \mu(U)}{2}$ then

$$(5.1) \quad (\mathcal{M}_m^\mu \downarrow U) \vee (\mathcal{M}_m^\mu \downarrow U) \subseteq \mathcal{M}_r^\mu \downarrow U.$$

Indeed, if O_1 and O_2 are elements of $\mathcal{M}_m^\mu \downarrow U$, we can assume that $\mu(O_1 \cup O_2) \leq \mu(U)$ so that

$$\mu(O_1 \cap O_2) = \mu(O_1) + \mu(O_2) - \mu(O_1 \cup O_2) > 2m - \mu(U) = r.$$

Self-splittability follows from the fact ⁽⁷⁾ that if $U_1 \cup U_2 \in \mathcal{M}_r$, for $d := \min(\mu(U_1), \mu(U_2), \mu(U_1 \cup U_2) - r) > 0$, $m_1 := \mu(U_1) - \frac{d}{2}$, $m_2 := \mu(U_2) - \frac{d}{2}$, we have

$$(5.2) \quad (\mathcal{M}_{m_1}^\mu \downarrow U_1) \cap (\mathcal{M}_{m_2}^\mu \downarrow U_2) \subseteq \mathcal{M}_r.$$

Indeed, if $A_i \in \mathcal{M}_{m_i}^\mu \downarrow U_i$ for $i \in \{1, 2\}$, then

$$\begin{aligned} \mu(A_1 \cup A_2) &= \mu(A_1) + \mu(A_2) - \mu(A_1 \cap A_2) \\ &\geq \mu(A_1) + \mu(A_2) - \mu(U_1 \cap U_2) \\ &> \mu(U_1) + \mu(U_2) - \mu(U_1 \cap U_2) - d \\ &> r. \end{aligned}$$

□

Theorem 5.2. *If X is a (Hausdorff) completely regular space but is not pre-Radon, then*

$$C_{\Lambda^\downarrow}(X, \mathbb{R}) > C_k(X, \mathbb{R}).$$

Proof. The proof of [2, Proposition 3.1] shows that if X is a Hausdorff non pre-Radon space, then there is a τ -additive finite measure μ and an $r > 0$ such that $\mathcal{M}_r$ is compact but not compactly generated. In view of Lemma 5.1, $\gamma_\mu \subseteq \Lambda^\downarrow(X)$ so that $C_k(X, \$^*) < C_{\Lambda^\downarrow}(X, \$^*)$. In view of Proposition 2.2, $C_k(X, \mathbb{R}) < C_{\Lambda^\downarrow}(X, \mathbb{R})$ because $\Lambda^\downarrow(X)$ is hereditary. □

Note that we have shown that the inclusion $k(X) \subseteq \Lambda^\downarrow(X)$ may be strict.

For instance, if X is the Sorgenfrey line, which is not pre-Radon, then $C_{\Lambda^\downarrow}(X, \mathbb{R})$ is a topological vector space and is strictly finer than the compact-open topology.

REFERENCES

- [1] B. Alleche and J. Calbrix, *On the coincidence of the upper Kuratowski topology with the cocompact topology*, Topology Appl., **93**(1999), 207-218.
- [2] A. Bouziad, *Borel measures in consonant spaces*, Topology Appl., **70** (1996), 125-132.
- [3] S. Dolecki, *Properties transfer between topologies on function spaces, hyperspaces and underlying spaces*, Mathematica Pannonica, **19**(2) (2008), 243-262.
- [4] S. Dolecki, G. H. Greco, and A. Lechicki, *When do the upper Kuratowski topology (homeomorphically, Scott topology) and the cocompact topology coincide?*, Trans. Amer. Math. Soc. **347** (1995), 2869-2884.
- [5] S. Dolecki and F. Mynard, *When is the Isbell topology a group topology?*, to appear in Topology Appl.
- [6] S. Dolecki and F. Mynard, *Relations between function spaces, hyperspaces and underlying spaces*, in preparation.
- [7] J.R. Isbell, *Function spaces and adjoints*, Math. Scandinavica **36** (1975), 317-339.
- [8] J.R. Isbell, *Meet-continuous lattices*, Symposia Mathematica **16** (1975), 41-54.
- [9] F. Jordan, *Coincidence of function space topologies*, submitted, 2008.
- [10] F. Jordan, *More on coincidence of function space topologies*, in preparation.
- [11] Elliot Pearl (ed.), *Open problems in topology II*, Elsevier, 2007.

⁷Note that γ_μ is not sectionable, so that Theorem 4.3 is not sufficient to deduce self-splittability.